

A close-up photograph of a honeybee on a bright yellow flower, likely a rapeseed flower, against a soft-focus background of more yellow flowers and a clear sky. The bee is positioned on the right side of the image, facing left towards the flower.

INTERNATIONAL APPEAL

Stop 5G on Earth and in Space

Sign the Appeal: www.5gSpaceAppeal.org

WIRELESS RADIATION AND OSTEOPOROSIS

I was astonished by the number of people who contacted me after I broke my arm telling me they had broken theirs too -- some of them this year, and others within the last few years. It occurred to me to wonder: has there been a significant increase in osteoporosis and bone fractures around the world? and if so, is this yet another health effect caused by the use of cell phones and their infrastructure irradiating our bones as well as the rest of our bodies?

I remembered reading some fascinating facts about bones in the groundbreaking 1985 book, *The Body Electric*, written by orthopedic surgeon Robert O. Becker. Bones, he discovered, are semiconductors, and they owe their electrical properties to being doped with tiny amounts of copper. The atoms of copper, he found, bond electrically to both apatite crystals and collagen fibers -- the two main components of bone -- and hold them together, "much as wooden pegs fastened the pieces of antique furniture to each other."

"Osteoporosis," wrote Becker, "comes about when copper is somehow removed from the bones. This might occur not only through chemical/metabolic processes, but by a change in the electromagnetic binding force, allowing the pegs to 'fall out.' It's possible that this could result from a change in the overall electrical fields throughout the body or from a change in those surrounding the body in the environment."

I also remembered, from the old Soviet Union literature, summarized in my 1997 book, [Microwaving Our Planet](#), that radio frequency radiation redistributes metals throughout the body.

With these facts in mind, I have searched the world's medical literature for studies on the incidence of both osteoporosis and fractures, and the evidence seems fairly conclusive: (1) There has been an enormous increase in the incidence of both osteoporosis and bone fractures of all types throughout the world in children and adults since about 1950; (2) the incidences of both continue to rise, worldwide; (3) most studies published in the past couple of decades have found that osteoporosis in children is correlated with the amount of time spent daily looking at screens; (4) rates of osteoporosis do *not* correlate with the amount of time children spend sitting but *not* looking at screens; and (5) these trends are independent of the amount of exercise people get.

The authors of these studies have been at a loss to explain their findings, but they are easily explained when one remembers the electrical properties of bones, and the effects that cell phone and computer screens, all emitting radiation, are likely to have on bones and on the copper atoms within them -- and that exposure to radiation from radio, TV, radar, and (more recently) cell tower antennas has increased tremendously since World War II.

Here is a sampling of the studies I have collected:

- Louis V. Avioli reviewed the world's literature in 1991. During the second half of the twentieth century, he found, both osteoporosis and fracture rates had risen dramatically in the United States, Canada, Norway, Sweden, Spain, Italy, the UK, Belgium, Australia, and elsewhere. The incidence rate of hip fractures in the United States had been increasing by about 40% per decade. (1)
- M.L. Grundill and M.C. Burger, in 2021, found that the incidence rate of hip fractures in a population in South Africa had more than doubled in men and almost sextupled in women compared to what had been reported in 1968. (2)
- Emmanuel K. Dretakis et al. found that the annual number of hip fractures in Crete increased 21% in just four years, from 1982 to 1986, while the population over 50 remained the same. (3)

- Hiroshi Koga et al. examined the records of children aged 6 to 14 in Niigata, Japan. The incidence rate of all fractures more than doubled from the early 1980s to the early 2000s in both girls and boys, and almost tripled in girls in junior high school. (4)
- P. Lühje et al. found that the incidence rate of hip fractures throughout Finland quadrupled between 1968 and 1988. (5)
- In 2012 Ambrish Mithal and Parjeet Kaur found that hip fracture rates had increased two- to three-fold throughout Asia during the previous 30 years. (6)
- Hiroshi Hagino et al. found that hip fracture rates in Tottori Prefecture, Japan had risen by almost 40% between 1986 and 1992, and by more than 60% in men and about 50% in women between 1986 and 2001. Increases in fracture rates occurred not only in the elderly, but in people in their 30s and 40s. (7)
- In 1989 Karl J. Obrant et al. did an analysis of fracture trends in Malmö, Sweden, where all X-rays have been saved since the beginning of the twentieth century. They found that the yearly number of fractures in that city had increased seven-fold between 1951 and 1985, and the incidence rate of fractures among children had doubled between 1950 and 1979. ***“There are signs that there is a deterioration of the quality of the skeleton in successive generations,”*** wrote the authors. ***“With the same or even diminished trauma, we sustain more serious and more comminuted fractures today than previously.”*** The increase had nothing to do with changing estrogen levels, because fracture rates had increased even more in men than in women. The daily consumption of both calcium and Vitamin D had increased during that time. But the incidence of hip fractures was higher in cities than in rural environments where, we know, there was less radiation. (8)
- Haiyu Shao et al., in 2015, looking at hours per day spent playing video games by Chinese adolescents, found that adolescents with longer video game time were more likely to have lower bone mass density in their legs, trunk, pelvis, spine, and whole body. (9)
- Anne Winther et al., studying 15- to 18-year-olds in Tromsø, Norway in 2010-2011, found that longer screen time was associated with lower bone mass density in both boys and girls, regardless of the amount of daily physical

activity, calcium intake, vitamin D, alcohol consumption, smoking habits, height or weight. (10)

- Sebastien Chastin, examining youths aged 8 to 22 in the U.S. in 2005-2006, found that screen-based sitting was associated with lower bone mass density in hips and spine. Non-screen-based sitting was *not* associated with lower bone mass density. (11)
- Natalie Lundin et al. found that annual incidence rates of pelvic and hip socket fractures in Sweden increased 25% from 2001 to 2016, and that increasing incidence rates were seen in all age groups. (12)
- Daniel Jerrhag et al. found that the incidence rate of forearm fractures in Sweden was 23% higher in 2010 compared with 1999, and that the increase was greater in men and women 17 to 64 years of age than in the elderly. (13)
- Michiel Herteleer et al. found that the incidence rate of pelvic and hip socket fractures in Belgium doubled between 1988 and 2006, and rose another 26% by 2018. (14)
- Neeraj M. Patel found that the annual incidence rate of fractures in children aged 6 to 18 in New York State almost quadrupled between 2006 and 2015. (15)

Donations to support our work are needed. The Cellular Phone Task Force is a 501(c)(3) nonprofit organization, and donations from U.S. residents are tax-deductible. Our Tax ID Number is 11-3394550.

DONATE

Arthur Firstenberg

Author, [*The Invisible Rainbow: A History of Electricity and Life*](#)

Administrator, [*International Appeal to Stop 5G on Earth and in Space*](#)

Caretaker, [*ECHOEarth \(End Cellphones Here On Earth\)*](#)

P.O. Box 6216

Santa Fe, NM 87502

USA

phone: +1 505-471-0129

info@cellphonetaskforce.org

May 11, 2022

The last 39 newsletters, including this one, are available for viewing on the [Newsletters page](#) of the Cellular Phone Task Force. Some of the newsletters are also available there in German, Spanish, Italian, French, Norwegian, and Dutch.

To subscribe, go to www.cellphonetaskforce.org/subscribe or click on this link:

SUBSCRIBE

References

- (1) Louis V. Avioli, "Significance of osteoporosis: A growing international health problem," *Calcified Tissue International* 49:S5-S7 (1991)
- (2) M.L. Grundill and M.C. Burger, "The incidence of fragility hip fractures in a subpopulation of South Africa," *South African Medical Journal* 111(9):896-902
- (3) Emmanuel K. Dretakis et al., "Increasing incidence of hip fracture in Crete," *Acta Orthopaedica Scandinavica* 63(2):150-151 (1992)
- (4) Hiroshi Koga et al., "Increasing incidence of fracture and its sex difference in school children: 20 year longitudinal study based on school health statistic in Japan," *Journal of Orthopaedic Science* 23(1):151-155 (2018)
- (5) P. Lüthje et al., "Increasing incidence of hip fracture in Finland," *Archives of Orthopaedic and Trauma Surgery* 112:280-282 (1993)
- (6) Ambrish Mithal and Parjeet Kaur, "Osteoporosis in Asia: A call to action," *Current Osteoporosis Reports* 10:245-247 (2012)
- (7) Hiroshi Hagino et al., "Increasing incidence of hip fracture in Tottori Prefecture, Japan: Trend from 1986 to 2001," *Osteoporosis International* 16:1963-1968 (2005)
- (8) Karl J. Obrant et al., "Increasing age-adjusted risk of fragility fractures," *Calcified Tissue International* 44:157-167 (1989)
- (9) Haiyu Shao et al., "Association between duration of playing video games and bone mineral density in Chinese adolescents," *Journal of Clinical Densitometry* 18(2):198-202 (2015)
- (10) Ann Winther et al., "Leisure time computer use and adolescent bone health -- findings from the Tromsø Study, *Fit Futures: a cross-sectional study*," *BMJ Open* 5:e006665 (2015)
- (11) Gadi Lissak, "Adverse physiological and psychological effects of screen time on children and adolescents: Literature review and case study," *Environmental Research* 164:149-157 (2018)

(12) Sebastian FM Chastin et al., "The frequency of osteogenic activities and the pattern of intermittence between periods of physical activity and sedentary behaviour affects bone mineral content: the cross-sectional NHANES study," *BMC Public Health* 14:4 (2014)

(13) Natalie Lundin et al., "Increasing incidence of pelvic and acetabular fractures. A nationwide study of 87,308 fractures over a 16-year period in Sweden," *Injury* 52:1410-1417 (2021)

(14) Daniel Jerrhag et al., "Epidemiology and time trends of distal forearm fractures in adults -- a study of 11.2 million person-years in Sweden," *BMC Musculoskeletal Disorders* 18, Article number 240 (2017)

(15) Michiel Herteleer et al., "Epidemiology and secular trends of pelvic fractures in Belgium: A retrospective, population-based, nationwide observational study," *Bone* 153:116141 (2021)